

CHAPTER 7

ARISTOLOCHIC ACID NEPHROPATHY: AN ENVIRONMENTAL AND IATROGENIC DISEASE

Arthur P. Grollman* and Bojan Jelaković†

Contents

1. Toxicity of Aristolochic Acid (AA)	218
2. Chemistry of AA	218
3. Chinese Herb Nephropathy (CHN)	219
4. Endemic (Balkan) Nephropathy (EN)	220
5. AAN = BEN = CHN	221
6. Dose–Toxicity Relationship	221
7. Renal Histopathology	222
8. Proximal Tubular Function	222
9. Urothelial (Transitional) Cell Cancer	223
10. AL-DNA Adducts	223
11. Mutational Signature of AA	225
12. Genetic Considerations	225
13. AAN as an Iatrogenic Disease	226
Acknowledgment	228
References	228

The *Aristolochiaceae* family of herbaceous plants, specifically members of the genus *Aristolochia*, has been used for medicinal purposes for more than 2500 years [1]. Remarkably, the extensive *materia medica* describing the therapeutic use of *Aristolochia* rarely mentions intrinsic toxicity. Recently, aristolochic acid (AA), a principal component of all *Aristolochia* sp., was shown to be the toxic principle responsible for the syndromes known as Chinese herb nephropathy (CHN) and endemic (Balkan) nephropathy (EN). Both disorders are associated with a high incidence of urothelial (transitional cell) cancer and appear to constitute a single disease entity, best designated as aristolochic acid nephropathy (AAN).

* Laboratory of Chemical Biology, Department of Pharmacological Sciences, Stony Brook University, Stony Brook, New York 11794

† Department of Nephrology and Arterial Hypertension, Zagreb University School of Medicine and University Hospital Center, 10000 Zagreb, Croatia, and Croatian Center for Endemic Nephropathy, 35000 Slavonski Brod, Croatia

The epidemiology and pathophysiology of CHN and EN have been reviewed extensively [2–7] and the association of these diseases with human cancer is the subject of several comprehensive reports [8,9]. Guided by the hypothesis that AA is the common etiologic agent in CHN and EN [10,11], we review the molecular and clinical toxicology of this powerful nephrotoxin–carcinogen. Additionally, based on the traditional use of *Aristolochia* in herbal remedies, we posit that AAN represents a long-overlooked iatrogenic disease and an international public health problem of considerable magnitude [12].

1. TOXICITY OF ARISTOLOCHIC ACID (AA)

Toxicologists have long been aware of the toxic properties of *Aristolochia* sp. In 1825, while investigating the homeopathic principle that “like cures like,” Jörg performed experiments on himself and his colleagues, revealing the acute toxic properties of *Aristolochia serpentaria* [13]. Among the effects experienced after ingesting multiple doses of the root (2.5–7.5 gm, estimated to contain ~5 mg AA) were intense gastrointestinal symptoms and polyuria. Later, Orfila, the founder of modern quantitative toxicology, demonstrated the lethal effects of *Aristolochia clematitis* in short-term experiments performed on dogs [14]. More than a century ago, Pohl administered the extracts prepared from the seeds and roots of *A. clematitis* to rabbits, revealing the nephrotoxic effects of the partially purified phytotoxin [15]. Dumić [16] and Martinčić [17] reported that horses ingesting hay contaminated with *A. clematitis* developed chronic renal failure and described the histopathology of the diseased kidneys [17]. Subsequently, Mengs documented the acute and chronic toxicities of purified AA in rodents, including its carcinogenicity [18–20]. The molecular and cellular mechanisms underlying AA cytotoxicity have been explored in cultured renal proximal tubule cells and in rodent models of AAN (reviewed in [21]).

2. CHEMISTRY OF AA

Aristolochia herbs contain a number of structurally related nitrophenanthrene carboxylic acids, principally aristolochic acid I (AA-I) and aristolochic acid II (AA-II) [22]. Many of the studies discussed here utilize the naturally occurring mixture of AA-I and AA-II designated for purposes of this discussion as AA.

3. CHINESE HERB NEPHROPATHY (CHN)

Between 1990 and 1992, ~1800 otherwise healthy Belgian women inadvertently ingested *Aristolochia fangchi* in conjunction with their participation in a weight-loss regimen. Several of these women developed rapidly progressive renal interstitial fibrosis leading to chronic renal failure [23]. Histopathologic examination revealed a corticomedullary gradient of interstitial fibrosis, and marked atrophy of proximal tubules with glomeruli generally being spared. Approximately 5% of women ingesting the herbal supplement developed end-stage renal disease (ESRD) ultimately requiring dialysis or renal transplantation [3]. When the diseased kidneys were removed, urothelial cell atypia and carcinoma were found in the upper urinary tract [24]. Specifically, among 39 patients undergoing prophylactic bilateral nephrectomy, 46% had urothelial cancers located primarily in the renal pelvis and upper ureter [25]. All but two of the remaining patients displayed mild-to-moderate urothelial cell dysplasia.

Definitive evidence that AA was responsible for CHN was obtained when aristolactam (AL)-DNA adducts were detected in renal tissues (reviewed in [26]). Additionally, Cosyns *et al.* reproduced the nephrotoxic and carcinogenic effects observed in humans by treating rabbits with low doses of AA [27]. Based on these reports, the term aristolochic acid nephropathy (AAN) was suggested to reflect the etiologic role of this phytotoxin in CHN [28].

The dramatic revelation that AA is a powerful nephrotoxin and carcinogen for humans drew attention to the worldwide distribution and use of *Aristolochia* sp. as herbal remedies. Not surprisingly, subsequent reports (reviewed in [9]) described almost 200 patients outside of Belgium in whom chronic renal failure followed ingestion of *Aristolochia* herbs.

Undoubtedly, published reports represent only a small fraction of AAN cases. In support of this assertion, we note that in a single year in China, 320,000 kg of *Aristolochia manchuensis* was harvested for medicinal purposes [29]. And, in 2003, 5000 kg of Qing Mu Xiang and 10,000 kg of Tian Xian Teng, *Aristolochia* herbals listed in the 2005 Chinese Pharmacopeia, were produced in the Shanghai district alone. Thus, despite heavy and widespread use of Aristolochic herbals in China, very few cases of AAN were reported prior to 1999 [30].

Epidemiologic studies in Taiwan, where the prevalence of chronic kidney disease (CKD) is among the highest in the world, reveal a strong association between CKD and the general use of herbal therapies [31]. Too, a survey of prescriptions filled between 1997 and 2003 in Taiwan revealed that one-third of the population had been treated with herbal products known to contain or suspected to be contaminated with *Aristolochia* [32].

CKD also is recognized as a major public health problem in India [33]. Traditional Ayurvedic medicines used in India and elsewhere include several species of *Aristolochia*, predominantly *Aristolochia indica* and *Aristolochia bracteolata*. Vanherweghem [34] has suggested that the nephrotoxicity associated with *Aristolochia* herbal remedies may contribute to the high incidence (27.8%) of chronic interstitial nephritis in India [35].

Thus, the failure to relate the traditional uses of *Aristolochia* sp. to its delayed nephrotoxic and carcinogenic effects has created a global health problem that will require the attention of public health authorities for the foreseeable future [13].

4. ENDEMIC (BALKAN) NEPHROPATHY (EN)

EN is a chronic tubulointerstitial disease occurring in rural villages located near tributaries of the river Danube in Bosnia and Herzegovina, Bulgaria, Croatia, Romania, and Serbia. First described 50 years ago [36–39], EN is estimated to affect at least 25,000 individuals in countries harboring this disease, while another 75,000 men and women residing in endemic regions are believed to be at risk [40]. Significant epidemiologic features of EN include its focal occurrence in certain farming villages; a familial but not inherited pattern of disease, frequently affecting members of the same household; occurrence only in individuals older than 18 years of age; presence in less than 10% of households in endemic villages; and a strong association with upper urinary tract transitional cell (urothelial) cancer (UUC) [4–6,41].

Until quite recently, the etiology of EN remained obscure [42]. A variety of environmental agents, including mycotoxins, heavy metals, viruses, and trace-element deficiencies have been investigated as potential causative agents [42,43], with ochratoxin A (OTA), being a principal focus of EN research in recent years. Residents of villages in endemic areas are exposed to relatively high concentrations of OTA; however, elevated serum levels of OTA have been detected in residents of countries throughout the world [44] while EN is limited to isolated villages in Balkan countries [4–6]. In fact, OTA has never been associated directly with nephrotoxicity in humans [45] and a comprehensive review of the literature concluded that published epidemiologic studies are inadequate to assess a cause-and-effect relationship between OTA and human cancer [46]. Thus, the weight of scientific evidence, including the inability of OTA to form DNA adducts under physiologic conditions [47,48], argues strongly against OTA or other mycotoxins playing a role in the etiology of EN [10,21].

Current experimental evidence strongly supports the view [49–51] that chronic dietary poisoning by AA accounts for all of the important

characteristics of EN, including the geographical distribution, characteristic pattern of renal tubulointerstitial fibrosis, and increased risk of developing UUC. Earlier, Kazantzis [52] and Ivić [53] suggested that *A. dematitidis* might contain the etiologic agent responsible for EN; however, this insightful observation was not pursued. Our impetus to investigate the role of AA in EN was prompted by the striking clinical and histopathologic similarities between EN and the nephropathy observed in the cluster of Belgian women [54].

5. AAN = BEN = CHN

In the following section, we review critically the toxicology of AA, focusing on renal proximal tubules and urothelial cells, which are targeted by the toxin. In so doing, we integrate observations on the pathophysiology of CHN and EN with research on animal models of AAN and mechanistic studies of the molecular and cellular effects of AA (reviewed in [21]).

6. DOSE-TOXICITY RELATIONSHIP

Estimates have been made of the amount of AA ingested by ~100 Belgian women who developed end-stage renal failure [3,55] and by many of the ~200 sporadic cases of CHN reported in the world literature (reviewed in [9]). Our analysis of the dose-toxicity relationship assumes that AA-induced DNA damage in cells is cumulative and largely irreversible. Thus, 1800 Belgian women received a maximum of 0.025 mg/kg of AA over a period averaging 13 months, with 5% of this group developing end-stage renal failure 3–85 months after administration was stopped [3]. Similarly, Chinese patients estimated to have ingested 0.7–1.5 mg AA per day intermittently over 1–10 years in the form of pills containing *A. mandhuriensis* developed chronic renal failure, sometimes within 6 months of initial exposure to the herb ([9] and X. Li, unpublished data). Higher doses of AA, such as those employed in a Phase I clinical study of AA (1 mg/kg for 3 or more days) [56], and in some Chinese patients (10 mg/day for 10 days) [57] resulted in acute renal toxicity.

In EN, 20 years or more may elapse before renal function becomes clinically apparent. The slow progression of this endemic disease likely reflects a low level of exposure to AA. Thus, Croatian farm families who ingest bread contaminated by AA are estimated to accumulate over 8–10 years the same amount of AA as did Belgium women with CHN in a single year [49]. The number of residents in Croatian endemic areas who depend on home-baked bread has decreased substantially in the past 20 years (Jelaković *et al.*, unpublished data). Decreased dietary exposure to AA is

expected to translate into a lower incidence of EN and a delayed age of onset of nephrotoxicity. The observed shift to the older age group among newly diagnosed EN cases and among EN patients beginning dialysis [5] is in line with this prediction.

7. RENAL HISTOPATHOLOGY

AAN is marked by progressive tubulointerstitial fibrosis resulting in ESRD. Comparison of the renal histopathology of patients with CHN and EN revealed a striking feature, namely, a corticomedullary gradient of interstitial fibrosis present in both diseases [54]. This gradient, reported only in AAN and cadmium-induced nephropathy [58] is even more pronounced in renal biopsies obtained prior to ESRD.

New Zealand White rabbits treated over a period of 17–21 months with low doses of AA developed impaired renal tubular function and interstitial fibrosis with a corticomedullary gradient [27]. Following treatment with AA, C3H/He mice developed acute tubular necrosis and interstitial fibrosis while C57BL/6 mice are resistant to similar doses of the toxin [59]. AA-I and AA-II exhibit similar genotoxic potential [60], however, AA-I is solely responsible for the observed nephrotoxicity in C3H mice [59,60]. Histopathologic findings in Wistar rats include necrosis of outer medullary proximal tubules and urothelial dysplasia. As in mice, renal injury in rats is manifested initially by an acute inflammatory/necrotic phase lasting several days, followed by a chronic phase, consisting of interstitial inflammation, tubular atrophy, and interstitial fibrosis [61]. Biochemical markers were consistent with epithelial–mesenchymal transition, though an alternative mechanism involving TGF- β activation of fibroblasts may account for some of the interstitial fibrosis involved [62].

8. PROXIMAL TUBULAR FUNCTION

AAN is characterized by proximal tubule dysfunction, including persistent glycosuria, aminoaciduria, and low-molecular weight (tubular) proteinuria (LMWP) [3,4]. Normally, LMWP are filtered at the glomerulus, reabsorbed by endocytosis in the proximal tubule, and catabolized. Proximal tubule damage interferes with this process; thus, the urinary LMWP/albumin ratio is strikingly elevated in patients with CHN and EN compared with patients with glomerular disease [63]. This parameter provides a useful biomarker for detecting early proximal tubule dysfunction in AAN [64–66].

Rabbits given AA intraperitoneally 5 days a week for 17–21 months developed impaired proximal tubule function manifested by glycosuria and

tubular proteinuria [27]. In Wistar rats, high doses of AA disrupted proximal tubule function, as reflected by proteinuria, glycosuria, and neutral amino acid enzymuria [67]. Both animal models reproduce the main features of AAN in humans. The rat was used to investigate early functional impairment of the proximal tubule, revealing that the outer medullary S3 segment of the proximal tubule was the preferential target of AA [67].

9. UROTHELIAL (TRANSITIONAL) CELL CANCER

The strong association between exposure to AA and the development of urothelial neoplasia was first recognized in Belgian women with CHN [24,68]. Ultimately, 40–45% of this group developed multifocal high-grade transitional cell carcinomas located primarily in the upper urinary tract [25]. Based on this observation, prophylactic bilateral nephroureterectomy has been strongly recommended for CHN and EN patients requiring dialysis or renal transplantation [69].

UUC also are prevalent among residents of endemic villages in the Balkan region. In the former Yugoslavia, data from more than 2000 cases of UUC were used to explore the epidemiology of this uncommon cancer [70]. The prevalence of UUC in certain endemic regions can be as much as 100-fold greater than in the country as a whole [71]. More than two-thirds of these cancers are localized in the renal pelvis or ureter [72,73] compared to ~5% of UUC among patients residing in nonendemic areas [74]. In Serbia, the fraction of urothelial cancers located in the renal pelvis and ureter increased from 18% in the period 1921–1952 to 63% in 1953–1960 [70,72].

Patients undergoing surgery for UUC often exhibit signs of renal dysfunction. Petković *et al.* recorded “renal failure” in 63% (111/187) of patients with UUC [75]. This important observation was confirmed by Nikolić in a review of 710 patients with UUC, 65% of whom had creatinine clearances of less than 80 ml/min [70]. Recently, 9 of 11 residents of endemic villages undergoing nephroureterectomy for UUC were found to have histopathologic changes in the renal cortex compatible with EN [50]. Thus, UUC may precede, accompany, or follow manifestations of nephrotoxicity in patients with EN. The age at which patients in endemic regions develop symptoms associated with UUC has increased steadily over the past 30 years [76] again suggesting that environmental exposure to AA is diminishing over time.

10. AL-DNA ADDUCTS

Both AA-I and AA-II are subject to enzymatic nitroreduction in mammalian cells, generating active nitrenium intermediates (Fig. 1) that react with the exocyclic amino groups of deoxyadenine and deoxyguanine




Figure 1 Principal urinary metabolites of aristolochic acids (AA) in rat. Cytochrome P450 (CYP) catalyzed demethylation of AA-I generates AA-Ia, which is subject, in turn, to Phase II biotransformation to form glucuronide, sulfate, and acetyl conjugates. Enzymatic nitroreduction (NR) of AA-I or AA-II yields the biologically inactive aristolactams AL-I and AL-II. Reactive nitrenium intermediates form covalent adducts with DNA and proteins.

bases of DNA (reviewed in [26]). Both AL-I- and AL-II-derived DNA adducts have been detected in the renal cortex of patients with BEN and/or UUC [50] and in persons ingesting botanical products containing AA (reviewed in [9]).

The level of AL-DNA adducts in target tissues represent the integration of external exposure with interindividual variability in metabolism, nucleotide excision repair, and other cellular processes, providing a tangible link between exposure to the genotoxin and its cytotoxic and carcinogenic effects [77]. Their usefulness as biomarkers is enhanced by their persistence in target tissues. For example, in humans, AL-DNA adducts were detected 36–89 months following termination of exposure to AA [26,50]. Thus, among Belgian women, AL-DNA adduct levels ranged from 0.1 to 17 adducts per 10^8 deoxynucleotides (dNs), with dA-AL-I being the major adduct found in the renal cortex while dA-AL-II and dG-AL-I were present in significantly lower amounts [26].

In patients with EN, adduct levels ranged from 8 to 59 adducts per 10^8 dNs for dA-AL-I and 2–62 adducts per 10^8 dNs for dG-AL-I [50]. In urothelial tumors, adduct levels were significantly lower, 0.7–1.6 and

0.3–0.5 adducts per 10^8 dNs for dA-AL and dG-AL, respectively [50]. In rodent models of AAN, AL-DNA adducts accumulate in target tissues [78]; however, the prominent cytotoxic effects of AA-I in proximal tubule cells appear unrelated to its DNA-damaging effects [60].

11. MUTATIONAL SIGNATURE OF AA

Mutational analysis of the p53 tumor suppressor gene in UUC tumors from patients residing in endemic villages revealed that the frequency of A:T \rightarrow T:A mutations was 78% [50]. In contrast, A:T \rightarrow T:A mutations occur infrequently in transitional cell carcinomas of the renal pelvis (0%), ureter (5%), and bladder (4.8%) in patients from nonendemic regions [79]. Importantly, the p53 mutation pattern in UUC associated with EN is consistent with the mutational spectra induced by AA-I (or by a mixture of AA-I and AA-II) in (i) the H-ras gene of rats [80]; (ii) transgenic rodent models [81]; (iii) the p53 gene in a urothelial cancer from a patient with CHN [82]; (iv) site-specific mutagenesis studies in which a single dA-AL adduct is transfected into NER-deficient human cells (Yang and Moriya, unpublished data); and (v) immortalized (Hupki) mouse cell lines carrying the human p53 gene [83]. Indeed, the predominance of A:T \rightarrow T:A transversions in the p53 mutational spectrum is widely recognized as a mutational signature for human exposure to AA [10,50,83,84].

12. GENETIC CONSIDERATIONS

Only 5% of the 1800 Belgian women exposed to similar amounts of AA developed AAN [3] and the reported prevalence of EN is 3–7% [5]. Similarly, various strains of mice exhibit differing susceptibilities to the toxic effects of AA [59,60]. Thus, genetically determined interindividual variability could contribute to the risk of developing AAN and/or its associated UUC [85]. Rarely, however, does a single mutated gene result in a high absolute risk of the associated cancer or other disease. Thus, susceptibility to AA most likely is influenced by the additive or synergistic effects of several genes, none of which individually may result in a large phenotypic effect. Furthermore, the cumulative effect of susceptibility to this environmental mutagen may result from complex interactions among multiple genes and from gene–environment interactions [86]. Molecular epidemiologic studies have been undertaken to potentially capture such interactive effects in EN, and mouse models of AAN have been used to identify genes controlling susceptibility to the nephrotoxic effects of AA [87].

Polymorphisms that affect susceptibility to AA are likely to be found in genes responsible for its absorption, bioactivation, detoxification, excretion, and/or transport. Metabolism of AA, mediated by enzymes known collectively as xenobiotic metabolizing enzymes (XMEs), has been demonstrated in humans and animals treated with AA [88] (Fig. 1). O-demethylation, mediated by Cytochrome P450s, results in rapid detoxification of AA-1 in rodents [89]. Thus, functional polymorphisms in XMEs may affect the level of critical intermediates produced during biotransformation, some of which bind to DNA and proteins. Similarly, polymorphisms in DNA repair enzymes can affect an individual's ability to excise the AL-DNA adducts that initiate UUC [90]. Although individual risks associated with such polymorphisms might be low, they have potentially great public health relevance (i.e., population-attributable risk) because of their high population frequency [91].

13. AAN AS AN IATROGENIC DISEASE

The varied uses of *Aristolochia* (birthwort) have been traced from the fourth century BC to the eighteenth century AD, based on excerpts from the original literature [1]. In a more specific approach, the actual cumulative doses of *Aristolochia* employed in treating illness during the Greco-Roman period were estimated by Scarborough who suggested that this early usage likely resulted in unrecognized iatrogenic disease as a result of the herb's intrinsic toxicity [92].

Extending this novel strategy, we have analyzed the use of *Aristolochia* herbal combinations in Ayurvedic and Chinese medicine. Our immediate goal is to document clearly the past and present use, worldwide, of *Aristolochia*, a family of herbs recently shown to be nephrotoxic and carcinogenic to humans.

The first description of *Aristolochia* is found in "Enquiry into Plants" by Theophrastus (300 BC) who described the morphology of the plant and its medical uses [94]. At that time, *Aristolochia* was used for snakebites, as a general antidote against poisons, to clean wounds, to alleviate insomnia, cure uterine malfunctions, ease bowel obstruction, and relieve symptoms of dropsy (edema). In the Hippocratic *Diseases of Women*, anonymous writers of that era discuss the use of *Aristolochia* as an oxytocic agent to aid women in childbirth and abortifacient, obstetric and gynecologic practices that continue to this day [95].

Arguably the most influential medical work of classical antiquity, *De Materia Medica*, written ca. 70 AD by Dioscorides, includes almost all herbs then employed in Greco-Roman pharmacology. Individual chapters are devoted to *Aristolochia rotunda*, *Aristolochia longa*, and *Aristolochia clematitis* [96]. Interestingly, not one word of caution was expressed regarding the potential toxicity of these herbs [92].

The influence of *De Materia Medica* on Western and Islamic medicine was profound. Dioscorides' descriptions of *Aristolochia* were translated into Arabic and Latin and this information was preserved throughout the Byzantine period and the Middle Ages [93]. With the advent of the printing press, information originating in *De Materia Medica* was incorporated (often verbatim) into the printed herbals of the Renaissance, which enjoyed wide distribution in Europe [97]. Ultimately, an entry for *Aristolochia* appeared in the *Pharmacopeia Londinensis of 1618* and; following that, in other official compendia and textbooks of pharmacology. Literature discussing properties of birthwort continued into the twentieth century [98]. Once again, usage of this herb was not associated with nephrotoxicity or cancer.

Ayurveda, one of several codified systems of traditional medicine, enjoys the same legal status in India as does allopathic medicine. The *Carakasamhita*, dating from ca. 400 AD and constituting one of the earliest Ayurvedic medical treatises, contains descriptions of 341 plant products. Scholars have identified in this classic text several species of *Aristolochia* used in Ayurveda today. Thus, patients currently treated by Ayurvedic practitioners are exposed to the toxic effects of these native herbs.

Gandhanakuli, *isvari*, and *nakuli*, identified as *Aristolochia indica* Linn or *A. bracteolata* Lam are mentioned in the *Carakasamhita* as antidotes to snakebite and for intermittent fevers (G.J. Meulenbeld, personal communication). Similar applications of *Aristolochia* sp. appear in more recent Indian pharmacopeia where the herb continues to be used for children with fever as well as an emmenagogue and antiarthritic [99].

In China, the earliest record of herbal medicines is the Shen Nong Ben Cao Jing (Shennong Herbal), compiled sometime between the first century BC and the second century AD [93]. Three hundred sixty-five agents were listed in this text and are divided into superior, medium, and inferior based on their relative toxicity [103]. Mou Dou Ling (fruit of *Aristolochia debilis*) first appears in Lei Gong Pao Jiu Lun (ca. 421 AD) with no mention of its toxic effects.

The Ben Cao Gang Mu, compiled by Li Shi-Zhen in the latter part of the sixteenth century, was based on the author's experience and on data obtained from earlier herbals. This Chinese herbal classic describes 1892 "drugs" (with 1110 drawings) including many species of *Aristolochia*. For 400 years, the Ben Cao Gang Mu remained the principal source of information in traditional Chinese medicine and the work was translated into numerous languages, reflecting its influence in countries other than China.

In the mid-twentieth century, Ben Cao Gang Mu was replaced by modern *materia medica*, the most comprehensive source being Zhong Hua Ben Cao (Encyclopedia of Chinese Materia Medica), published in 1999 [100]. The Encyclopedia lists 23 species of *Aristolochia*, again with little mention of toxicity. The Chinese government currently lists the following *Aristolochia* herbs: *A. manshuriensis* (stems), *A. fangchi* (root), *A. debilis* (root)

and fruit), and *A. contorta* (*fruit*), two of which (Mou Dou Ling and Quingmuxiang) appear in the 2005 Pharmacopeia.

For thousands of years, herbs have been used for medicinal purposes and in the twenty-first century 80% of the world's population still rely on herbal remedies to treat symptoms of disease. *Aristolochia* herbals play an important role in this tradition. The governments of China and India, countries that each have populations in excess of one billion, have been slow in implementing measures to reduce human exposure to this toxic herb. Thus, from a global perspective, we conclude that AAN and its associated upper urothelial cancer continues to exist as it has in centuries past as a "silent" but omnipresent iatrogenic disease.

ACKNOWLEDGMENT

The authors express their appreciation to Professor John Scarborough, Professor Gerrit Jan Meulenbeld and Ms. Jie Wang, respectively, for their translations of ancient Greek, Sanscrit and Chinese text. We thank Dr. Jovan Nikolic for his critique of the section on upper urinary tract cancer and Ms Annette Oestreicher for her skilled editorial work. The National Institute of Environmental Health Sciences (grant ES-04068), Fogarty International Center (grant 1203-07042) and Croatian Ministry of Science supported our collaborative research on aristolochic acid nephropathy.

REFERENCES

- [1] W.R. Dawson, Birthwort: A study of the progress of medical botany through twenty-two centuries, *Pharmaceutical J. Pharmacist* 396–397 (1927) 427–430.
- [2] J.P. Cosyns, Aristolochic acid and "Chinese herb nephropathy": A review of the evidence to date, *Drug Safety* 26 (2003) 33–48.
- [3] J.L. Vanherweghem, F. DeBelle, M.C. Muniz-Martinez, J. Nortier, Aristolochic acid nephropathy after Chinese herb remedies, In: M.E. De Broe, G.A. Porter, W.M. Bennett, and G.A. Verpooten (Eds.), *Clinical Nephrotoxins* 2nd Edition, Kluwer, Dordrecht, 2003, pp. 579–603.
- [4] L. Djukanović, Z. Radovanović, in: M.E. De Broe, G.A. Porter, W.M. Bennett, and G.A. Verpooten (Eds.), *Balkan Endemic Nephropathy in Clinical Nephrotoxins* 2nd Edition, Kluwer, Dordrecht, 2003, pp. 588–601.
- [5] D. Bukvic, Endemic nephropathy, overview of investigations in the Lazarevac area, Professional Pen Co, Belgrade, 2004.
- [6] V. Stefanović, Z. Radovanović, Z. Balkan, Endemic nephropathy and associated urothelial cancer, *Nat. Clin. Pract. Urol.* 5 (2008) 105–112.
- [7] G. Bamias, J. Boletis, Balkan nephropathy: Evolution of our knowledge, *Am. J. Kidney Dis.* 52 (2008) 606–616.
- [8] *Aristolochia* species and aristolochic acids, In IARC Monographs on the evaluation of the carcinogenic risk of chemicals to humans: Some traditional herbal medicines, some mycotoxins, naphthalene and styrene Vol. 82, pp. 69–128, International Agency for Research on Cancer, Lyon, France, 2002.

- 1 [9] Aristolochic Acids, In National Toxicology Program Report on Carcinogens 12th 1
- 2 Edition, US Dept Health and Human Services, Public Health Service, Research 2
- 3 Triangle Park, North Carolina, 2008. 3
- 4 [10] A.P. Grollman, B. Jelaković, Role of environmental toxins in endemic (Balkan) 4
- 5 nephropathy, J. Am. Soc. Nephrol. 18 (2007) 2817–2823. 5
- 6 [11] H. de Jonge, Y. Vanrenterghem, Aristolochic acid: The common culprit of 6
- 7 Chinese herbs nephropathy and Balkan endemic nephropathy, Nephrol. Dial. 7
- 8 Transplant. 23 (2007) 39–41. 8
- 9 [12] F.D. Debelle, J.L. Vanherweghem, J.L. Nortier, Aristolochic acid nephropathy: 9
- 10 A worldwide problem, Kidney Int. 74 (2008) 158–169. 10
- 11 [13] J.C.G. Jörg, Materialien zu einer künftigen Heilmittellehre durch Versuche der 11
- 12 Arzneien an gesunden Menschen gewonnen und gesammelt von Dr. Johann 12
- 13 Christian Gottfried Jörg, Carl Cnobloch, Leibzig, 1825. 13
- 14 [14] M.J.B. Orfila, de l'Aristolochie (*Aristolochia clematidis*), In Traité. des poisons tiré des 14
- 15 régnés minéral, végétal et animal; ou, Toxicologie générale, considère sous les rapports de 15
- 16 la physiologie et de la pathologie et de la médecine légale, 1851, pp. 296–298. 16
- 17 [15] J. Pohl, Ueber das Aristolochin, einen giftigen Bestandtheil der *Aristolochia*arten, Arch. 17
- 18 Exp. Path. u. Pharm. 29 (1892) 282–302. 18
- 19 [16] A. Dumić, Trovanje konja vučjom stopom (*A. clematidis* L), Izdanje vojnotehničkog 19
- 20 glasnika, Belgrade, Yugoslavia. 1954, pp. 3–35. 20
- 21 [17] M. Martinčić, Toxische einwirkungen der *Aristolochia clematidis* auf die Niere des 21
- 22 Pferdes, VeterinarSKI Arhiv. 27 (1958) 51–59. 22
- 23 [18] U. Mengs, W. Lang, J.A. Poch, The carcinogenic action of aristolochic acid in rats, 23
- 24 Arch. Toxicol. 51 (1982) 107–119. 24
- 25 [19] U. Mengs, Acute toxicity of aristolochic acid in rodents, Arch. Toxicol. 59 (1987) 25
- 26 328–331. 26
- 27 [20] U. Mengs, C.D. Stotzem, Toxicity of aristolochic acid—a subacute study in male rats, 27
- 28 Med. Sci. Res. 20 (1992) 223–224. 28
- 29 [21] K.G. Dickman, A.P. Grollman, Nephrotoxicity of natural products: Aristolochic acid 29
- 30 and fungal toxins, in: R.G. Schnellman (Ed.), Comprehensive Toxicology: Renal 30
- 31 Toxicology 2nd Edition, Elsevier, New York, 2009. 31
- 32 [22] V. Kumar, A.K. Prasad, V.S. Parmar, Naturally occurring aristolactams, aristolochic 32
- 33 acids and diisoaporphines and their biological activities, Nat. Prod. Rep. 20 (2003) 33
- 34 565–583. 34
- 35 [23] J.L. Vanherweghem, M. Depierreux, C. Tielemans, D. Abramowicz, M. Dratwa, 35
- 36 M. Jadoul, *et al.*, Rapidly progressive interstitial renal fibrosis in young women: 36
- 37 Association with slimming regimen including Chinese herbs, Lancet 341 (1993) 37
- 38 387–391. 38
- 39 [24] J.P. Cosyns, M. Jadoul, J.P. Squifflet, P.J. van Cangh, C. van Ypersele de Strihou, 39
- 40 Urothelial malignancy in nephropathy due to Chinese herbs, Lancet 344 (1994) 188. 40
- 41 [25] J.L. Nortier, M.C. Martinez, H.H. Schmeiser, V.M. Arlt, C.A. Bieler, M. Petein, 41
- 42 *et al.*, Urothelial carcinoma associated with the use of a Chinese herb (*Aristolochia* 42
- 43 *fangchi*), N. Engl. J. Med 342 (2000) 1686–1692. 43
- 44 [26] V.M. Arlt, M. Stiborová, H.H. Schmeiser, Aristolochic acid as a probable human 44
- cancer hazard in herbal remedies: A review, Mutagenesis 17 (2002) 265–277. 45
- [27] J.P. Cosyns, J.P. Dehoux, Y. Guiot, *et al.*, Chronic aristolochic acid toxicity in rabbits: 46
- A model of Chinese herb nephropathy?, Kidney Int. 5 (2001) 2164–2173. 47
- [28] G. Gillerot, M. Jadoul, V.M. Arlt, *et al.*, Aristolochic acid nephropathy in a Chinese 48
- patient: Time to abandon the term “Chinese herbs nephropathy”? Am. J. Kidney Dis. 49
- 38 (2001) E26. 50
- [29] S.L. Hu, H.Q. Zhang, K. Chan, Q.X. Mei, Studies on the toxicity of *Aristolochia* 51
- manshuriensis* (Guanmuton), Toxicology 198 (2004) 195–201. 52

- 1 [30] X. Li, H. Wang, Aristolochic acid nephropathy: What we know and what we have to 1
do, *Nephrology* 9 (2004) 109–111. 2
- 3 [31] J.Y. Guh, H.C. Chen, J.F. Tsai, L.Y. Chuang, Herbal therapy is associated with the 3
risk of CKD in adults not using analgesics in Taiwan, *Am. J. Kidney Dis.* 49 (2007) 4
626–633. 4
- 5 [32] S.C. Hsieh, I.H. Lin, W.L. Tseng, C.H. Lee, J.D. Wang, Prescription profile of 5
potentially aristolochic acid containing Chinese herbal products: An analysis 6
of National Health Insurance data in Taiwan between 1997 and 2003, *Chin. Med.* 7
3 (2008) 13. 7
- 8 [33] G.K. Modi, V. Jha, The incidence of end-stage renal disease in India: A population- 8
based study, *Kidney Int.* 70 (2006) 2131–2133. 9
- 10 [34] J.L. Vanherweghem, *Aristolochia* sp. and chronic interstitial nephropathies in Indians, 10
Lancet 349 (1997) 1399. 11
- 12 [35] M.K. Mani, Chronic renal failure in India, *Nephrol. Dial. Transplant.* 8 (1993) 12
684–689. 13
- 14 [36] Y. Tanchev, Z. Evstatiev, D. Dorossiev, J. Pencheva, G. Zvetkov, Studies of the 14
nephritides in the District of Vratza, *Savrememna Medicina* 7 (1956) 14–29. 15
- 16 [37] V. Danilović, M. Djurišić, M. Mokranjac, B. Stojimirović, J. Živoljinović, 16
P. Stojaković, Porodična oboljenja bubrega u selu Šopić izazvana hroničnom intok- 17
sikacijom olovom, *Srpski Arh Lek* 85 (1957) 1115–1125. 17
- 18 [38] N. Foarta, M. Negoescu, Nefrita cronica azotemia endo-epidemic Stud, *Cercet* 18
Med. 1 (1961) 217–221. 19
- 19 [39] World Health Organization, Memorandum: The endemic nephropathy of South- 19
Eastern Europe, *Bull. World Health Organ.* 32 (1965) 441–448. 20
- 21 [40] C.A. Tatu, W.H. Orem, R.B. Finkelman, G.L. Feder, The etiology of Balkan 21
endemic nephropathy: Still more questions than answers, *Environ. Health Perspect.* 22
106 (1998) 689–700. 23
- 24 [41] S. Čeović, M. Miletić-Medved, Epidemiological features of endemic nephropathy in 24
the focal area of Brodska Posvina, Croatia, in: D. Čvorišćec, S. Čeović, and 25
A. Stavljenić-Rukavina (Eds.), *Endemic Nephropathy in Croatia*, 1996, pp. 7–21. 25
- 26 [42] V. Batuman, Fifty years of Balkan endemic nephropathy: Daunting questions, elusive 26
answers, *Kidney Int.* 69 (2006) 644–646. 27
- 28 [43] T.C. Voice, D.T. Long, Z. Radovanović, J.L. Atkins, *et al.*, Critical evaluation of 28
environmental exposure agents suspected in the etiology of Balkan endemic nephrop- 29
athy, *Int. J. Occup. Environ. Health* 12 (2006) 369–376. 29
- 30 [44] H.A. Clark, S.M. Snedeker, A. Ochratoxin, Its cancer risk and potential for exposure, 30
J. Toxicol. Environ. Health B Crit. Rev. 3 (2006) 265–296. 31
- 32 [45] M. Godin, J.P. Fillastre, A. Fancois, F. Le Roy, J.P. Morin, Is Ochratoxin A 32
nephrotoxic in human beings? *Adv. Nephrol.* 26 (1997) 181–206. 33
- 34 [46] J. Fink-Gremmels, A. Ochratoxin, In food: Recent developments and significance, 34
Food Addit. Contam. 22(Suppl. 1), (2005) 1–5. 35
- 35 [47] A. Mally, H. Zepnik, P. Wanek, E. Eder, E.K. Dingley, H. Ihmels, W. Volkel, 35
W. Dekant, Ochratoxin A: Lack of formation of covalent DNA adducts, *Chem. Res.* 36
Toxicol. 17 (2004) 234–242. 37
- 38 [48] T. Delatour, A. Mally, J. Richoz, S. Özden, W. Dekant, H. Ihmels, D. Otto, 38
D. Gasparutto, M. Marin-Kuan, B. Schilter, C. Cavin, Absence of 2'-deoxyguano- 39
sine-carbon-8-bound ochratoxin A adduct in rat kidney DNA monitored by isotope 40
dilution LC-MS/MS, *Mol. Nutr. Food Res.* 52 (2008) 472–482. 40
- 41 [49] T. Hranjec, A. Kovač, J. Kos, W. Mao, J.J. Chen, A.P. Grollman, B. Jelaković, 41
Endemic nephropathy: The case for chronic poisoning by aristolochia, *Croat Med J.* 42
46 (2005) 116–125. 43
- 44 44

- [50] A.P. Grollman, S. Shibutani, M. Moriya, F. Miller, L. Wu, U. Moll, *et al.*, Aristolochic acid and the etiology of endemic (Balkan) nephropathy, *Proc. Natl. Acad. Sci. USA* 104 (2007) 12129–12134.
- [51] H. de Jonge, Y. Vanrenterghem, Aristolochic acid: The common culprit of Chinese herbs nephropathy and Balkan endemic nephropathy, *Nephrol. Dial. Transplant.* 23 (2008) 39–41.
- [52] G. Kazantzis, Comment in the general discussion of possible nephrotoxic agents, in: G.E. Wolstenholme and J. Knight (Eds.), *The Balkan Nephropathy*, CIBA Foundation, Boston, MA, 1967, p. 114.
- [53] M. Ivić, The problem of etiology of endemic nephropathy, *Liječ Vjes* 91 (1969) 1273–1281.
- [54] J.P. Cosyns, M. Jadoul, J.P. Squifflet, J.F. de Plaen, D. Ferluga, C. van Ypersele de Strihou, Chinese herbs nephropathy: A clue to Balkan endemic nephropathy? *Kidney Int.* 45 (1994) 1680–1688.
- [55] M.C.M. Martinez, J. Nortier, P. Vereerstraeten, J.L. Vanherweghem, Progression rate of Chinese herb nephropathy: impact of *Aristolochia fangchi* ingested dose, *Nephrol. Dial. Transplant.* 17 (2002) 408–412.
- [56] L. Jackson, S. Kofman, A. Weiss, H. Brodovsky, Aristolochic acid (NSC-50413): phase I clinical study, *Cancer Chemother. Rep.* 42 (1964) 35–37.
- [57] S.H. Lo, K.L. Mo, K.S. Wong, S.P. Poon, C.K. Chan, C.K. Lai, A. Chan, Aristolochic acid nephropathy complicating a patient with focal segmental glomerulosclerosis, *Nephrol. Dial. Transplant.* 19 (2004) 1913–1915.
- [58] M. Yasuda, A. Miwa, M. Kitagawa, Morphometric studies of renal lesions in Itai-Itai disease: chronic cadmium nephropathy, *Nephron* 69 (1995) 14–19.
- [59] N. Sato, D. Takahashi, S.M. Chen, R. Tsuchiya, T. Mukoyama, S. Yamagata, *et al.*, Acute nephrotoxicity of aristolochic acids in mice, *J. Pharm. Pharmacol.* 56 (2004) 221–229.
- [60] S. Shibutani, H. Dong, N. Suzuki, S. Ueda, F. Miller, F.A.P. Grollman, Selective toxicity of aristolochic acids I and II, *Drug Metab. Dispos.* 35 (2007) 1217–1222.
- [61] F.D. Debelle, J.L. Nortier, E.G. de Prez, C.H. Garbar, A.R. Vienne, I.J. Salmon, M.M. Deschodt-Lanckman, J.L. Vanherweghem, Aristolochic acids induce chronic renal failure with interstitial fibrosis in salt-depleted rats, *J. Am. Soc. Nephrol.* 13 (2002) 431–436.
- [62] A.A. Pozdzik, I.J. Salmon, F.D. Debelle, C. Decaestecker, C.C. van den Branden, D. Verbeelen, M.M. Deschodt-Lanckman, J.L. Vanherweghem, J.L. Nortier, Aristolochic acid induces proximal tubule apoptosis and epithelial to mesenchymal transformation, *Kidney Int.* 73 (2008) 595–607.
- [63] A. Kabanda, M. Jadoul, R. Lauwerys, A. Bernard, C. van Ypersele de Strihou, Low molecular weight proteinuria in Chinese herbs nephropathy, *Kidney Int.* 48 (1995) 1571–1576.
- [64] P.W. Hall, M. Piscator, M. Vasiljević, N. Popović, Renal function studies in individuals with the tubular proteinuria of endemic Balkan nephropathy, *Q. J. Med.* 41 (1972) 385–393.
- [65] M. Radonić, Beta-2 microglobulin in Balkan endemic nephropathy, *Pathol. Biol.* 26 (1978) 317–320.
- [66] D. Čvorišćec, Early diagnosis of endemic nephropathy, *Clin. Chim. Acta* 297 (2000) 85–91.
- [67] C. Lebeau, F.D. Debelle, V.M. Arlt, A. Pozdzik, E.G. De Prez, D.H. Phillips, M.M. Deschodt-Lanckman, J.L. Vanherweghem, J.L. Nortier, Early proximal tubule injury in experimental aristolochic acid nephropathy: functional and histological studies, *Nephrol. Dial. Transplant.* 20 (2005) 2321–2332.

- 1 [68] J.P. Cosyns, M. Jadoul, J.P. Squifflet, F.X. Wese, C. van Ypersele de Strihou, Urothelial lesions in Chinese-herb nephropathy, *Am. J. Kidney Dis.* 33 (1999) 1011–1017.
- 2 [69] B. Jelakovic, *et al.*, Consensus statement of the World Health Organization collaborative Center for primary health care: international workshop on screening, diagnosis, classification and treatment of endemic (Balkan) nephropathy.
- 3 [70] J. Nikolić, Epidemic Nephropathy and Upper Urothelial Tumors, Izdavačko preduzeće, Belgrade, Serbia, 2006.
- 4 [71] S. Petković, Epidemiology and treatment of renal pelvic and ureteral tumors, *J. Urol.* 114 (1975) 858–865.
- 5 [72] V. Petronić, Tumors of the upper urothelium and endemic nephropathy, in: Z.Rado-
vanovic, M. Sindic, M. Polenakovic, L. Djukanović, and V. Petronic (Eds.), Endemic
6 Nephropathy, Zavod Za Udzbenike I Nastavna Sredstva, Belgrade Serbia, 2000,
pp. 350–439.
- 7 [73] M. Miletić-Medved, A.M. Domijan, M. Peraica, Recent data on endemic nephropathy and related urothelial tumors in Croatia, *Wien Klin. Wochenschr.* 117 (2005) 604–609.
- 8 [74] E.M. Genega, C.R. Porter, Urothelial neoplasm of the kidney and ureter. An epidemiologic, pathologic, and clinical review, *Am. J. Clin. Pathol.* 117(Suppl.), (2002) S36–S48.
- 9 [75] S. Petković, Correlation of endemic nephropathy and tumors of renal pelvic and ureter, In First Congress of Nephrologists of Yugoslavia, Belgrade, 1981, pp. 177–182.
- 10 [76] J. Nikolić, M. Djokić, I. Ignjatović, V. Stefanović, Upper urothelial tumors in emigrants from Balkan endemic nephropathy areas in Serbia, *Urol. Int.* 77 (2006) 240–244.
- 11 [77] G.N. Wogan, S.S. Hecht, J.S. Felton, A.H. Conney, L.A. Loeb, Environmental and chemical carcinogenesis, *Sem. Cancer Biol.* 473 (2004) 486.
- 12 [78] H. Dong, N. Suzuki, M.C. Torres, R.R. Bonala, F. Johnson, A.P. Grollman, S. Shibutani, Quantitative determination of aristolochic acid-derived DNA adducts in rats using ³²P-postlabeling/polyacrylamide gel electrophoresis analysis, *Drug Metab. Dispos.* 34 (2006) 1122–1127.
- 13 [79] M. Olivier, R. Eeles, M. Hollstein, M.A. Khan, C.C. Harris, P. Hainaut, The IARC TP53 database: new online mutation analysis and recommendations to users, *Hum Mutat.* 19 (2002) 607–614.
- 14 [80] H.H. Schmeiser, J.W. Janssen, J. Lyons, H.R. Scherf, W. Pfau, A. Buchmann, C.R. Bartram, M. Wiessler, Aristolochic acid activates *ras* genes in rat tumors at deoxyadenosine residues, *Cancer Res.* 50 (1990) 5464–5469.
- 15 [81] A. Kohara, T. Suzuki, M. Honma, T. Ohwada, M. Hayashi, Mutagenicity of aristolochic acid in the lambda/lacZ transgenic mouse (mutatmmouse), *Mutat. Res.* 515 (2002) 63–72.
- 16 [82] G.M. Lord, T. Cook, V.M. Arlt, H.H. Schmeiser, G. Williams, C.D. Pusey, Urothelial malignant disease and Chinese herbal nephropathy, *Lancet* 358 (2001) 1515–1516.
- 17 [83] T. Nedelko, V.M. Arlt, D.H. Phillips, M. Hollstein, TP53 mutation signature supports involvement of aristolochic acid in the etiology of endemic nephropathy-associated tumors, *Int. J. Cancer* 124 (2008) 987–990.
- 18 [84] V.M. Arlt, M. Stiborová, J. vom Brocke, M.L. Simões, G.M. Lord, J.L. Nortier, M. Hollstein, D.H. Phillips, H.H. Schmeiser, Aristolochic acid mutagenesis: Molecular clues to the aetiology of Balkan endemic nephropathy-associated urothelial cancer, *Carcinogenesis* 28 (2007) 2253–2261.
- 19 [85] R. Their, K. Golka, T. Bruning, *et al.*, Genetic susceptibility to environmental toxicants; the interface between human and experimental studies in the development of new toxicological concepts, *Toxicol. Lett.* 127 (2002) 321–327.

- [86] S.N. Kelada, D.L. Eaton, W.W. Wang, N.R. Rothman, M.J. Khoury, The role of genetic polymorphisms in environmental health, *Environ. Health Perspect.* 111 (2003) 1055–1064.
- [87] T.A. Rosenquist, P.A. Strockbine, A. Smith, H. Einolf, A.P. Grollman, Genetics of aristolochic acid nephropathy in mice, *Cold Spring Harb.* (2008).
- [88] G. Krumbiegel, J. Hallensleben, W.H. Mennicke, N. Rittman, H.J. Roth, Studies on the metabolism of aristolochic acids I and II, *Xenobiotica* 17 (1987) 981–991.
- [89] S. Shibutani, R.R. Bonala, T. Rosenquist, R. Rieger, N. Suzuki, F. Johnson, F. Miller, A.P. Grollman, Detoxification of aristolochic acid-I by O-demethylation: Less nephrotoxicity and genotoxicity of aristolochic acid Ia in rodents, *Carcinogenesis*, 2008.
- [90] E.C. Friedberg, G.C. Walker, W. Seide, R.D. Wood, R.A. Schultz, T. Ellenberger, *DNA Repair and Mutagenesis*, ASM Press, Washington, DC, 2006.
- [91] N. Caporaso, A. Goldstein, Cancer genes—single and susceptibility—exposing the difference, *Pharmacogenetics* 5 (1995) 59–63.
- [92] J. Scarborough, Kidney disease among the Romans, *Coll. Antropol.* 30(Suppl. 1) (2006).
- [93] R.H. Major, *A History of Medicine*, C.C Thomas, Springfield, Ill, 1954.
- [94] J. Scarborough, Theophrastus on herbals and herbal remedies, *J. Hist. Biol.* 11 (1978) 353–358.
- [95] J. M. Riddle, *Eve's Herbs, A History of Contraception and Abortion in the West*, Harvard University Press, Cambridge, MA, 1997.
- [96] Dioscorides, *Materia Medica*, III, 4, in: M. Wellman (Ed.), *Pedanii Dioscuridis Anazarbei De material medica*, 3 vols, Weidmann, Berlin, 1906–1914 (rptd. 1958; Vol. 2, pp. 6–8).
- [97] A. Arber, *Herbals: Their Origin and Evolution, A Chapter in the History of Botany 1470–1670* 3rd Edition, Cambridge Univ Press, Cambridge, .
- [98] M. Grieve, *A modern herbal: The medicinal, culinary, cosmetic and economic properties, cultivation and folk-lore of herbs, grasses, fungi, shrubs and trees with all their modern scientific uses*, Dover, New York, 1971, p. 104.
- [99] G. Watt, *A Dictionary of the Economic Products of India*, Vol. 1, 331–336, Superintendent of government printing, Calcutta, 1885.
- [100] Zhong Hua Ben Cao, 3–460–509 edited by the Health Department and National Chinese Medicine Management Office, published by Shanghai Science Technology Publication, 1999.

Au7

Au8

UNCORRECTED PROOF

Author Query Form

Book Series: Advances in Molecular Toxicology, 3
Chapter 07



Dear Author,

During the preparation of your manuscript for typesetting some questions have arisen. These are listed below. Please check your typeset proof carefully and mark any corrections in the margin of the proof or compile them as a separate list. This form should then be returned with your marked proof/list of corrections to Elsevier Science.

Disk use

In some instances we may be unable to process the electronic file of your article and/or artwork. In that case we have, for efficiency reasons, proceeded by using the hard copy of your manuscript. If this is the case the reasons are indicated below:

- ☐ Disk damaged ☐ Incompatible file format ☐ LaTeX file for non-LaTeX journal
- ☐ Virus infected ☐ Discrepancies between electronic file and (peer-reviewed, therefore definitive) hard copy.
- ☐ Other:

We have proceeded as follows:

- ☐ Manuscript scanned ☐ Manuscript keyed in ☐ Artwork scanned
- ☐ Files only partly used (parts processed differently:.....)

Bibliography

If discrepancies were noted between the literature list and the text references, the following may apply:

- ☐ The references listed below were noted in the text but appear to be missing from your literature list. Please complete the list or remove the references from the text.
- ☐ Uncited references: This section comprises references which occur in the reference list but not in the body of the text. Please position each reference in the text or, alternatively, delete it. Any reference not dealt with will be retained in this section.

Query Refs.	Details Required	Author's response
AU1	Please check and confirm for Introduction part.	
AU2	Please expand the term "NER" at its first occurrence if it is an acronym.	
AU3	This reference is not provided in the reference list. Please add to list or delete from text.	
AU4	Please provide the contribution title for this reference.	
AU5	Please list out all the author names in all such cases.	
AU6	Please provide complete bibliographic details for this reference.	
AU7	Please provide the volume and page number.	
AU8	Check page range.	